## WASHINGTON STATE UNIVERSITY



## PACIFIC NORTHWEST EXTENSION **ACTIVITIES ON JOINTED GOATGRASS**

## Joseph P. Yenish\*, Roland Schirman, **Doug Schmale, and Eric Zakarison,** Washington State University, Pullman

The Steering Committee of the National Jointed Goatgrass Research Program recognized that effective technology transfer was critical to the success of the program. Thus, the National Extension Coordinator Position was developed in the



The current arrangement of two part-time extension coordinators in different locations, both administered through WSU, has been successful. The potential for collaboration between programs has increased while still allowing extension efforts to better address regional needs. Dr. Schirman and Mr. Schmale both have experience in technology transfer. More importantly, their experience in winter wheat production systems in their respective regions better allows for directed programming and delivery. Both individuals have expressed a willingness in providing informational updates for the website on a voluntary basis once the program has formally ended. WSU-Extension has been the publication agency for all Extension Bulletins (Figure 2). These efforts will continue through two additional bulletins. Centralizing publication production has provided a consistency of format which adds to program recognition. WSU's Department of Crop and Soil Sciences also provides technical web assistance. The departmental communications coordinator is responsible for updating website information. Again this technical resource is under the same WSU administrative unit as the extension coordinators, but in much closer physical proximity. This closeness has streamlined the website update process. This centralization will aid in the expansion of the website to include additional grass weed species (Figure 3).

initial phase of the program.

The first two individuals to hold the position; Dr. Brian Jenks, University of Nebraska, and Dr. Mack Thompson, **Colorado State University; made a priority of increasing** awareness of the jointed goatgrass problem among farmers and grain handlers. The third coordinator, Eric Zakarison, Washington State University (WSU), took information more directly to growers via popular press articles, direct contact, and further expansion of the website. Dr. Tony White, Kansas State University, continued direct grower contact, expanded use of the media, and even further developed the website. Following Dr. White's departure, the current system of two regional extension coordinators, Dr. Roland Schirman for the Intermountain and Pacific Northwest Regions along with **Doug Schmale for the Southern and Central Grain Plains** Regions developed. Currently, both Roland's and Doug's positions are administered by WSU Extension although both work from their respective homes in Washington and Nebraska.

As the program matured, WSU became the center of technology transfer for the National Jointed Goatgrass Research Program. In addition to employing the two extension coordinators, WSU houses the webpage (Figure 1) has published and distributed extension bulletins generated by the program (Figure 2).

<text><text><text><text><text><text><text><text><text><text></text></text></text></text></text></text></text></text></text></text>	To a second of the	
<text><text><text><text><text><text><text></text></text></text></text></text></text></text>	production by severely reducing yields and	
Agglops cylindrica (Age-uh-lops sa-lin-dri-ca), the scientific name for jointed goatgrass. However, the impact this weed has on winter wheat producers big dollars. Domestic and export customers of U.S. wheat are currently demanding low dockage grain. Grain high in dockage is difficult to sell, expensive to ransport, and requires additional cleaning. Fai- ure by the United States to export clean wheat oran makes it easier for foreign competitors to increase their own market share to free orgen competitors to increase having lower profit potential (and greature producers) having lower profit potential (and greature producers) toon risk) than winter wheat. Jointed goatgrass in pro- having lower profit potential (and greature producers) heaving lower profit potential (and greature producers) seeds can persist in the soil for many years. The use of a 3- or 4 year crop rotations. Bradlead or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat haddition, growers must integrate alternate crops into theri normal rotation when jointed goatgrass infesta- tions are sever. The introduction of Clearfield <sup>4</sup>		
<ul> <li>scientific name for jointed goatgrass. However, the impact this weed has on winter wheat producers, the winder this weed has on winter wheat producers of the set of control of jointed goatgrass costs wheat producers big dollars.</li> <li>Domestic and export customers of U.S. wheat are currently demanding low dockage grain. Grain high in dockage is difficult to sell, expensive to transport, and requires additional clean wheat or module of jointed goatgrass. Research Program we haite a twow.jointedgoatgrass.org.</li> <li>Domestic needs have forceign competitors to increase their own market share in the world wheat market.</li> <li>Dense infestations of jointed goatgrass in production fields have forceig growers to abandorstandard rotations and switch to spring croppaying lower profit potential (and greater production risk) than winter wheat. Jointed goatgrass requires an integrated goatgrass control.</li> <li>Minimizing economic losses from jointed goatgrass control.</li> <li>Minimizing economic losses from jointed goatgrass control.</li> <li>Minimizing economic losses from jointed goatgrass infestions are sever. The introduction of Cleanfield?</li> <li>Domestic neeses are theretate alternate crops into their origin rotations when integrate alternate crops into their origin rotations when the soutions. In addition, growers must integrate alternate crops into their origin rotations when integrate alternate crops into their origin rotations when integrate alternate crops into their origin rotations of cleanfield?</li> </ul>		
ers gues far beyond issues of enunciation. Jointed goatgraas costs wheat producers big dollars. Domestic and export customers of U.S. wheat are currently demanding low dockage grain. Grain high in dockage is difficult to sell, expensive to anakes it easier for foreign competitors to increase their own market share in the world wheat market beir own market share in the world wheat market shandard rotations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations of jointed goatgrass is pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3 - or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadlead rot warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management trategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are sever. The introduction of Cleanfield <sup>1</sup>	scientific name for jointed goatgrass. However,	have prepared a series of management bulletins
goargrass costs wheat producers big dollars. Domestic and export customers of U.S. wheat are interpreted by the United States. They also will be available on the National Jointed Goargrass Research Program website at www.jointedgoargrass.org. Control of jointed goargrass from scientists across the United States. They also will be available on the National Jointed Goargrass Research Program website it a www.jointedgoargrass.org. Control of Jointed goargrass from scientists across the United States. They also will be available on the National Jointed Goargrass Research Program website it a www.jointedgoargrass.org. Control of Jointed goargrass research Program website at www.jointedgoargrass.org. Control for the National Jointed goargrass across the United States. They also will be available on the National Jointed goargrass.org. Control Program website at www.jointedgoargrass.org. Control in winter wheat. United goargrass in production fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater production fields have forced growers to allow tillage before spring planting and the use of various selective herbicides for jointed goargrass control. Minimizing economic losses from jointed goargrass control. In minimizing profitable crop rotations. In addition, growers must integrate alternate crops into their formal rotation when jointed goargrass infestations are severe. The introduction of Clearfield <sup>10</sup>		
Domestic and export customers of U.S. wheat are currently demanding low dockage grain. Grain high in dockage is difficult to sell, expensive to transport, and requires additional cleaning. Fai- ure by the United States to export clean wheat only makes it easier for foreign competitors to increase their own market share in the world wheat market. Dense infestrations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a <sup>3</sup> - or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbark populations to more acceptable levels. Broadlead or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement strategies for growing winter wheat while main- tining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>30</sup>		
Domestic and export customers of U.S. wheat are currently demanding low dockage grain. Grain high in dockage is difficult to sell, expensive to transport, and requires additional cleaning. Failure the bythe United States to export clean wheat onty makes it easier for foreign competitors to increase their own market share in the world wheat market. Dense infestations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation will winter wheat may be necessary to reduce soil seedBank opulations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, gowers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>9</sup>	goargrass costs wheat producers big douars.	
currently demanding low dockage grain. Grain high in dockage is difficult to sell, expensive to transport, and requires additional cleaning. Failure we by the United States to export clean wheat ondor makes it easier for foreign competitors to increase their own market share in the world wheat market. Dense infestations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadlead for warm-season grass crops can allow tilage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>a</sup>	Domestic and export customers of U.S. wheat are	사실 전에 가장 전에 가장 것 같아요. 이렇게 가장 이 것 같아요. 이렇게 다 가장 가지 않는 것이 같아요. 이렇게 나와 가지 않는 것이 나라 가지 않는 것이 같아요. 이렇게 나라 가지 않는 것이 나라 가지 않는 것이 같아요. 이렇게 나라 가지 않는 것이 같아요. 이렇게 다 가지 않는 것이 같아요. 이렇게 아니 않는 것이 같아요. 이렇게 않는 것이 같아요. 이렇게 아니 않는 것이 같아요. 이렇게 아니 않
<section-header><text><text><text><text></text></text></text></text></section-header>	The second s	1. And the product of the product
<section-header><text><text><text><text></text></text></text></text></section-header>	high in dockage is difficult to sell, expensive to	
makes it easier for foreign competitors to increase their own market share in the world wheat market. Dense infestations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbark populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>9</sup>		
Hakes it easier for foreign competitors to increase their own market share in the world wheat market. Dense infestations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while maint- nining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield?		Control
Dense infestations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3 - or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbark populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield?		
Dense infestations of jointed goatgrass in pro- duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3 - or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield?	ucu own maner snare in the world wheat market.	Historically jointed maternes has been difficult
duction fields have forced growers to abandon standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>a</sup>	Dense infestations of jointed goatgrass in pro-	
standard rotations and switch to spring crops having lower profit potential (and greater produc- tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>9</sup>		
tion risk) than winter wheat. Jointed goatgrass seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield®		ACCURATE AND A CONTRACTOR OF A CONTRACTOR OF A
seeds can persist in the soil for many years. The use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass control. Minimizing economic losses from jointed goatgrass control strategies for growing winter wheat while maintaining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infestations are severe. The introduction of Clearfield*		1 11 41 11 5/10
use of a 3- or 4-year crop rotation with winter wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield*		
wheat may be necessary to reduce soil seedbank populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield*		
populations to more acceptable levels. Broadleaf or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>a</sup>		
or warm-season grass crops can allow tillage before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>a</sup>		
before spring planting and the use of various selective herbicides for jointed goatgrass control. Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield®		The second s
Minimizing economic losses from jointed goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield®	before spring planting and the use of various	A REAL REAL PLAN AND A REAL
goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>®</sup>	selective herbicides for jointed goatgrass control.	
goatgrass requires an integrated approach by producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>®</sup>	Material and a second state of the second stat	STATES IN THE UPACE IN THE REAL PROPERTY IN
producers. They must implement management strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield®		
strategies for growing winter wheat while main- taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>®</sup> Jointed goatgrass infesting winter wheat.		THE ALL REAL PROPERTY OF A DECK AND
taining profitable crop rotations. In addition, growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield <sup>a</sup> Jointed goatgrass infesting winter wheat.		A CONTRACTOR OF A CONTRACT
growers must integrate alternate crops into their normal rotation when jointed goatgrass infesta- tions are severe. The introduction of Clearfield® Jointed goatgrass infesting winter wheat.		
tions are severe. The introduction of Clearfield® Jointed goatgrass infesting winter wheat.		
		and show a second second second second second
www.jointedgoatgrass.org	tions are severe. The introduction of Clearfield®	Jointed goatgrass infesting winter wheat.
		www.jointedgoatgrass.org





	tgrass Research Program - Windows Internet Explorer			
💽 🗸 😽 ktp://joir	ntedgoatgrass.wsu.edu/jointedgoatgrass/index.htm	🖌 🛃 🗙	Google	
File Edit View Favor	ites Tools Help			
🚱 🔓 National Joint	ed Goatgrass Research Program	🕴 - 📾 - 🖶	🔹 🔂 Page 🔹 🎯 Tools 🔹 🕢 🚽	3
	Grass Weeds In Wheat	eturn to Grass Weeds In Wheat		
Jointed Goatgrass Home	Jointed Goatgrass	Information Contacts		
NJGGRP	Now available!	Doug Schmale		
Photo Gallery		Extension Coordinator (Great Plains) National JGG Research Program		
	ER ///// Lountod (-ostarsee Host Manadomont)			
Identification &	<ul> <li>EB2003 - Jointed Goatgrass, Best Management Practices, Intermountain Region</li> </ul>	Lodgepole, NE 308-483-5080		
Biology	Practices, Intermountain Region	308-483-5080		
Biology Genetics				
Biology Genetics Management	Practices, Intermountain Region	308-483-5080 schmale@wsu.edu States Covered by Doug:		
Biology Genetics Management Bulletins	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico,		
Biology Genetics Management Bulletins Other Publications	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug:		
Biology Genetics Management Bulletins	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed.</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado, Nebraska, Wyoming		
Biology Genetics Management Bulletins Other Publications Contacts	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed. There have been multiple times and sites of introduction of JGG</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado,		
Biology Genetics Management Bulletins Other Publications Contacts Links	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed. There have been multiple times and sites of introduction of JGG into North America. Some speculate that it was transported into Kansas around the late 19th century in contaminated 'Turkey'</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado, Nebraska, Wyoming Roland Schirman Extension Coordinator (Intermountain/PNW)		
Biology Genetics Management Bulletins Other Publications Contacts	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed. There have been multiple times and sites of introduction of JGG into North America. Some speculate that it was transported into Kansas around the late 19th century in contaminated 'Turkey' wheat seed brought from Russia by Mennonite settlers. Others</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado, Nebraska, Wyoming Roland Schirman Extension Coordinator		
Biology Genetics Management Bulletins Other Publications Contacts Links	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed. There have been multiple times and sites of introduction of JGG into North America. Some speculate that it was transported into Kansas around the late 19th century in contaminated 'Turkey'</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado, Nebraska, Wyoming Roland Schirman Extension Coordinator (Intermountain/PNW) National JGG Research Program		
Biology Genetics Management Bulletins Other Publications Contacts Links	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed. There have been multiple times and sites of introduction of JGG into North America. Some speculate that it was transported into Kansas around the late 19th century in contaminated Turkey' wheat seed brought from Russia by Mennonite settlers. Others reported it was brought into the Kansas in wheat made by the</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado, Nebraska, Wyoming Roland Schirman Extension Coordinator (Intermountain/PNW) National JGG Research Program Dayton, WA (509) 382-2778 roland@wsu.edu		
Biology Genetics Management Bulletins Other Publications Contacts Links	<ul> <li>Practices, Intermountain Region</li> <li>EB2033E - Jointed Goatgrass, Best Management Practices, Central Great Plains</li> <li>Jointed goatgrass (Aegilops cylindrica L.) is a winter annual grass native to Asia and Europe. The weed was thought to be introduced into the U.S. as a contaminant in winter wheat seed. There have been multiple times and sites of introduction of JGG into North America. Some speculate that it was transported into Kansas around the late 19th century in contaminated 'Turkey' wheat seed brought from Russia by Mennonite settlers. Others reported it was brought into the Kansas in wheat made by the USDA, private seed firms, or individuals during the early 1900s.</li> </ul>	308-483-5080 schmale@wsu.edu States Covered by Doug: Texas, New Mexico, Oklahoma, Kansas, Colorado, Nebraska, Wyoming Roland Schirman Extension Coordinator (Intermountain/PNW) National JGG Research Program Dayton, WA (509) 382-2778		

## Grass Weeds In Wheat - Windows Internet Explorer 😋 🔿 👻 🍓 http://jointedgoatgrass.wsu.edu/index.htm 🗸 😽 🗙 Google File Edit View Favorites Tools Hel 🖓 🔹 🔝 🕘 🛲 🔹 🔂 Page 🔹 🚳 Tools 👻 🔞 👯 🤌 🙀 🍄 😽 Grass Weeds In Wheat **Grass Weeds** Monday, March 2, 2009 Your Grass ointed Goatgrass Downy Brome Rattail Fescue Searc Resource Welcome to GRASS WEEDS IN WHEAT!! This site is an expansion of work completed under the National Jointed Goatgrass Research Program which was funded by series of CSREES-USDA special grants from 1994 - 2009. For information and directed links on problem grass weeds, select from the grass species of interest to you. ontact us: yenish@wsu.edu , 509-335-2961 | Accessibility | Copyright | Policies pe Yenish, Extension Weed Scientist, Washington State University, PO Box 646420, Pullman, WA, 99164-6420 USA Grass Weeds In Wheat - Copyright © 2008 Disclaimer | Nondiscrimination Statemen Design & Hosting Provided by Washington State University Extension Figure 3. The National Jointed Goatgrass Research Program is

expanding to include additional winter annual grass weeds.

Figure 1. The homepage of the National Jointed Goatgrass Research Program. (www.jointedgoatgrass.org) is currently under expansion to include additional winter annual grass weeds.

**Development of the website was an initial technology** transfer priority of the program with instructions to develop on-line information geared towards producer usage. Following its origins in Nebraska the website moved into the Kansas State University System, and recently to the computer network of WSU. Work is underway to expand the website to include other winter annual grass weeds troublesome in wheat.

> Figure 2. Extension bulletins are published by Washington State University and are available as downloadable PDF files.

Winter wheat growers in the western states infested with jointed goatgrass have benefited from the diversity and efforts of all individuals who have served in the role of extension coordinator. As the program concludes, WSU will continue to house the jointedgoatgrass.org website expanding it into a central source of information on annual grass weeds in winter wheat. Such efforts will require continued collaboration of weed scientists through organizations such as the WERA-077, Managing Invasive Weeds in Wheat. The National Jointed Goatgrass Research Program has provided the base for continued collaborative efforts on grass weeds in wheat.